Author Index

Aanesrud, A., see Rasmussen, K.E. 355
Adachi, T., see Mogi, M. 194
Alary, J.
—, Carrera, G., De Saint Blanquat, G., Anglade, F. and Escrieut, C.
Reversed-phase ion-pair high-performance liquid chromatographic determination of 2-carboxythiazolidine-4-carboxylic acid in plasma 485
Angelo, H.R.
—, Herrstedt, J. and Jørgensen, M.
High-performance liquid chromatographic method with fluorescence detection for the simultaneous determination of metopimazine and its acid metabolite in serum 472
Anglade, F., see Alary, J. 485
Aoyama, T., see Ohtsubo, K. 397
Aramaki, S., see Tanaka, T. 407
Ashmore, S.P.
—, Thomson, A.H. and Simpson, H.
High-performance liquid chromatographic technique for the rapid determination of histamine in both the plasma and cellular components in blood 435
Ayers, C.R., see Herold, D.A. 180
Baldew, G.S.
—, De Goeij, J.J.M. and Vermeulen, N.P.E.
Determination of ^{75}Se-labelled selenite and metabolites in plasma and urine by high-performance liquid chromatography with on-line radioactivity detection 111
Bannwarth, B., see Lapicque, F. 301
Beeman, C.S.
— and Rossomando, E.F.
Assay of ornithine decarboxylase activity by reversed-phase high-performance liquid chromatography 101
Bever, C.T., see Leslie, J. 214
Bhaumik, K.
— and Huisman, T.H.J.
Application of high-performance liquid chromatographic methodology to the analysis of hemoglobins synthesized in erythroid progenitor cells 83
Bhavnagri, V.P., see Osman, M.A. 478
Blanckaert, N., see Heirwegh, K.P.M. 1
Bree, H. de, see Lammers, N. 291
Brubaker, Jr., W.F.
— and Williams, E.D.
Rapid, high-sensitivity reversed-phase liquid chromatographic assay for 9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazolin-5-imine and its oxo metabolite in plasma using fluorescence detection 377
Bulbena, O., see Ramis, I. 416
Carrera, G., see Alary, J. 485
Carver, A.H., see Ocampo, A.P. 167
Chamart, S.
—, Hanocq, M., Helson, M., Devleeschouwer, N. and Leclercq, G.
Determination of 2-methyl derivatives of tamoxifen in cell culture medium using high-performance liquid chromatography and electrochemical detection 365
Chan, Y.-P.M.
—, Lee, F.-W. and Siu, T.-S.S.
Quantitation of homoharringtonine in plasma by high-performance liquid chromatography with amperometric detection 155
Chanon, M., see Saleh, S. 223
Cheng, L.K., see Osman, M.A. 478
Davidsson, P.
—, Fredman, P. and Svennerholm, L.
Gangliosides and sulphatide in human cerebrospinal fluid: quantitation with immunoaffinity techniques 279
De Bree, H., see Lammers, N. 291
De Goeij, J.J.M., see Baldew, G.S. 111
De Haan, A., see Van Doorn, J.E. 441
De Jong, B.J., see Lammers, N. 291
De Saint Blanquat, G., see Alary, J. 485
Descombe, J.-J.
—, Doumont, G. and Picard, M.
Determination of exifone in human plasma and urine by high-performance liquid chromatography with electrochemical detection 345
Devleeschouwer, N., see Chamart, S. 365
Doorn, J.E. van, see Van Doorn, J.E. 441
Doumont, G., see Descombe, J.-J. 345
Dugan, M.E.R
—, Sauer, W.C. and Robinson, P.H.
Cellulose clean-up and high-performance liquid chromatography of DL-diaminopi-

Author Index

Aanesrud, A., see Rasmussen, K.E. 355
Adachi, T., see Mogi, M. 194
Alary, J.
—, Carrera, G., De Saint Blanquat, G., Anglade, F. and Escrieut, C.
Reversed-phase ion-pair high-performance liquid chromatographic determination of 2-carboxythiazolidine-4-carboxylic acid in plasma 485
Angelo, H.R.
—, Herrstedt, J. and Jørgensen, M.
High-performance liquid chromatographic method with fluorescence detection for the simultaneous determination of metopimazine and its acid metabolite in serum 472
Anglade, F., see Alary, J. 485
Aoyama, T., see Ohtsubo, K. 397
Aramaki, S., see Tanaka, T. 407
Ashmore, S.P.
—, Thomson, A.H. and Simpson, H.
High-performance liquid chromatographic technique for the rapid determination of histamine in both the plasma and cellular components in blood 435
Ayers, C.R., see Herold, D.A. 180
Baldew, G.S.
—, De Goeij, J.J.M. and Vermeulen, N.P.E.
Determination of ^{75}Se-labelled selenite and metabolites in plasma and urine by high-performance liquid chromatography with on-line radioactivity detection 111
Bannwarth, B., see Lapicque, F. 301
Beeman, C.S.
— and Rossomando, E.F.
Assay of ornithine decarboxylase activity by reversed-phase high-performance liquid chromatography 101
Bever, C.T., see Leslie, J. 214
Bhaumik, K.
— and Huisman, T.H.J.
Application of high-performance liquid chromatographic methodology to the analysis of hemoglobins synthesized in erythroid progenitor cells 83
Bhavnagri, V.P., see Osman, M.A. 478
Blanckaert, N., see Heirwegh, K.P.M. 1
Bree, H. de, see Lammers, N. 291
Brubaker, Jr., W.F.
— and Williams, E.D.
Rapid, high-sensitivity reversed-phase liquid chromatographic assay for 9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazolin-5-imine and its oxo metabolite in plasma using fluorescence detection 377
Bulbena, O., see Ramis, I. 416
Carrera, G., see Alary, J. 485
Carver, A.H., see Ocampo, A.P. 167
Chamart, S.
—, Hanocq, M., Helson, M., Devleeschouwer, N. and Leclercq, G.
Determination of 2-methyl derivatives of tamoxifen in cell culture medium using high-performance liquid chromatography and electrochemical detection 365
Chan, Y.-P.M.
—, Lee, F.-W. and Siu, T.-S.S.
Quantitation of homoharringtonine in plasma by high-performance liquid chromatography with amperometric detection 155
Chanon, M., see Saleh, S. 223
Cheng, L.K., see Osman, M.A. 478
Davidsson, P.
—, Fredman, P. and Svennerholm, L.
Gangliosides and sulphatide in human cerebrospinal fluid: quantitation with immunoaffinity techniques 279
De Bree, H., see Lammers, N. 291
De Goeij, J.J.M., see Baldew, G.S. 111
De Haan, A., see Van Doorn, J.E. 441
De Jong, B.J., see Lammers, N. 291
De Saint Blanquat, G., see Alary, J. 485
Descombe, J.-J.
—, Doumont, G. and Picard, M.
Determination of exifone in human plasma and urine by high-performance liquid chromatography with electrochemical detection 345
Devleeschouwer, N., see Chamart, S. 365
Doorn, J.E. van, see Van Doorn, J.E. 441
Doumont, G., see Descombe, J.-J. 345
Dugan, M.E.R
—, Sauer, W.C. and Robinson, P.H.
Cellulose clean-up and high-performance liquid chromatography of DL-diaminopi-

melic acid in hydrolysates of physiological samples 430
Dunning, L.K., see Osman, M.A. 478
Duyn, P. van, see Van Welie, R.T.H. 463
Eadie, M.J., see Herkes, G.K. 147
Embree, L.
— and McErlane, K.M.
Development of a high-performance liquid chromatographic-post-column fluorogenic assay for digoxin in serum 321
Escrieut, C., see Alary, J. 485
Felder, R.A., see Kinter, M. 201
Fevery, J., see Heirwegh, K.P.M. 1
Fredman, P., see Davidsson, P. 279
Fujii, N., see Ohtsubo, K. 397
Gelpi, E., see Ramis, I. 416
Gerlo, E., see Sion, J.-P. 91
Gillet, P., see Lapicque, F. 301
Ginneken, C.A.M. van, see Russel, F.G.M. 234
Goeij, J.J.M. de, see Baldew, G.S. 111
Goormachtig, P.F.J., see Van Doorn, J.E. 441
Gorus, F., see Sion, J.-P. 91
Goto, I., see Ohtsubo, K. 397
Gribnau, F.W.J., see Russel, F.G.M. 234
Groen, C.P., see Lammers, N. 291
Haan, A. de, see Van Doorn, J.E. 441
Hanocq, M., see Chamart, S. 365
Harada, M., see Mogi, M. 194
Hartmann, C.
—, Krauss, D., Spahn, H. and Mutschler, E.
Simultaneous determination of (*R*)- and (*S*)-celiprolol in human plasma and urine: high-performance liquid chromatographic assay on a chiral stationary phase with fluorimetric detection 387
Heirwegh, K.P.M.
—, Fevery, J. and Blanckaert, N.
Review. Chromatographic analysis and structure determination of biliverdins and bilirubins 1
Helson, M., see Chamart, S. 365
Henry, J.A., see Virgili, P. 228
Herkes, G.K.
—, McKinnon, G.E. and Eadie, M.J.
Simultaneous quantitation of salivary carbamazepine, carbamazepine-10,11-epoxide, phenytoin and phenobarbitone by high-performance liquid chromatography 147
Herold, D.A.
—, Smithy, B.J., Ross, R.M., Marquis, F., Savory, J., Wills, M.R. and Ayers, C.R.
Measurement of plasma thromboxane B_2 by gas chromatography–mass spectrometry using $^{18}O_2$-labelled thromboxane B_2 as the internal standard 180
Herrstedt, J., see Angelo, H.R. 472
Higuchi, S., see Ohtsubo, K. 397
Hoyt, K.D., see Ocampo, A.P. 167
Huisman, T.H.J., see Bhaumik, K. 83
Joerg, E.A., see Tremaine, L.M. 423
Johnston, A., see Saleh, S. 223
Jong, B.J. de, see Lammers, N. 291
Jørgensen, M., see Angelo, H.R. 472
Justová, V.
Thin-layer chromatography as an efficient alternative to high-performance liquid chromatography in the assay of plasma vitamin D_3 metabolites 242
Kato, H., see Nagata, O. 456
Kawahara, Y., see Shioya, H. 129
Kežić, S.
—, Kuiper, S. and Monster, A.C.
Gas chromatographic determination of 2,3-butanediol isomers in urine 209
Kinter, M.
—, Singh, S. and Felder, R.A.
Quantitation of selective dopaminergic drugs in plasma by gas chromatography–mass spectrometry following solid-phase extraction 201
Kiriyama, N., see Nagata, O. 456
Kojima, K., see Mogi, M. 194
Krauss, D., see Hartmann, C. 387
Krcmarik, C. Sanders, see Sanders Krcmarik, C. 493
Krehbiel, D.K., see Mifune, M. 55
Kuiper, S., see Kežić, S. 209
Kuroki, S., see Yamashita, H. 255
Lambert, H., see Lapicque, F. 301
Lammers, N.
—, De Bree, H., Groen, C.P., Ruijten, H.M. and De Jong, B.J.
Determination of drug protein-binding by high-performance liquid chromatography using a chemically bonded bovine albumin stationary phase 291
Lant, M.S., see Oxford, J. 137
Lapicque, F.
—, Netter, P., Bannwarth, B., Trechot, P., Gillet, P., Lambert, H. and Royer, R.J.
Identification and simultaneous determi-

nation of non-steroidal anti-inflammatory drugs using high-performance liquid chromatography 301
Laureys, M., see Sion, J.-P. 91
Leclercq, G., see Chamart, S. 365
Lee, F.-W., see Chan, Y.-P.M. 155
Leslie, J.
— and Bever, C.T.
Analysis of 3,4-diaminopyridine in human serum by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 214
Levy, R.J., see Sintov, A. 335
McErlane, K.M., see Embree, L. 321
McKinnon, G.E., see Herkes, G.K. 147
Marquis, F., see Herold, D.A. 180
Martin, D.S., see Sawyer, R.C. 450
Matsui, H.
— and Sekiya, T.
High-performance liquid chromatographic separation of urinary hippuric and *o*-, *m*- and *p*-methylhippuric acids with a *β*-cyclodextrin-bonded column 189
Meijel, J.J.M. van, see Russel, F.G.M. 234
Miceli, J.J., see Sanders Krcmarik, C. 493
Mifune, M.
—, Krehbiel, D.K., Stobaugh, J.F. and Riley, C.M.
Multi-dimensional high-performance liquid chromatography of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Mogi, M.
—, Harada, M., Adachi, T., Kojima, K. and Nagatsu, T.
Selective removal of *β*2-microglobulin from human plasma by high-performance immunoaffinity chromatography 194
Momose, A., see Tanake, T. 407
Moncrieff, J.
Determination of pharmacological levels of harmane, harmine and harmaline in mammalian brain tissue, cerebrospinal fluid and plasma by high-performance liquid chromatography with fluorimetric detection 269
Monster, A.C., see Kežić, S. 209
Murata, K.
—, Yokoyama, Y. and Yoshida, K.
High-performance liquid chromatographic identification of disaccharides generated from heparan sulphate isomers using heparitinases 27
Mutschler, E., see Hartmann, C. 387
Nagao, T.
— and Tanimura, T.
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using high-performance liquid chromatography with amperometric detection 39
Nagata, O.
—, Sakashita, T., Takahara, E., Kato, H. and Kiriyama, N.
Determination of a new muscle relaxant (HY-770) in human serum by gas chromatography–mass spectrometry 456
Nagatsu, T., see Mogi, M. 194
Nakayama, F., see Yamashita, H. 255
Netter, P., see Lapicque, F. 301
Ocampo, A.P.
—, Hoyt, K.D., Wadgaonkar, N., Carver, A.H. and Puglisi, C.V.
Determination of tazobactam and piperacillin in human plasma, serum, bile and urine by gradient elution reversed-phase high-performance liquid chromatography 167
O'Hare, M., see Smith, J.A. 71
Ohtsubo, K.
—, Higuchi, S., Aoyama, T., Fujii, N. and Goto, I.
Sensitive determination of ambenonium chloride in serum from patients with myasthenia gravis using ion-exchange resin extraction and reversed-phase ion-pair chromatography 397
Osman, M.A.
—, Dunning, L.K., Bhavnagri, V.P. and Cheng, L.K.
Determination of muzolimine in plasma and urine by high-performance liquid chromatography 478
Oxford, J.
— and Lant, M.S.
Development and validation of a liquid chromatographic–mass spectrometric as-

say for the determination of sumatriptan in plasma 137
Pachla, L.A., see Sanders Krcmarik, C. 493
Picado, C., see Ramis, I. 416
Picard, M., see Descombe, J.-J. 345
Puglisi, C.V., see Ocampo, A.P. 167
Ramis, I.
—, Roselló-Carafau, J., Bulbena, O., Picado, C. and Gelpi, E.
15-Hydroxyeicosatetraenoic acid as a major eicosanoid in nasal secretions: assay by high-performance liquid chromatographic–radioimmunoassay and gas chromatographic–mass spectrometric procedures 416
Rasmussen, K.E.
—, Tønnesen, F., Thanh, H.H., Rogstad, A. and Aanesrud, A.
Solid-phase extraction high-performance liquid chromatographic determination of flumequine and oxolinic acid in salmon plasma 355
Riley, C.M., see Mifune, M. 55
Robinson, P.H., see Dugan, M.E.R. 430
Rogstad, A., see Rasmussen, K.E. 355
Roselló-Carafau, J., see Ramis, I. 416
Ross, R.M., see Herold, D.A. 180
Rossomando, E.F., see Beeman, C.S. 101
Royer, R.J., see Lapicque, F. 301
Ruijten, H.M., see Lammers, N. 291
Russel, F.G.M.
—, Tan, Y., Van Meijel, J.J.M., Gribnau, F.W.J. and Van Ginneken, C.A.M.
Solid-phase extraction of furosemide from plasma and urine and subsequent analysis by high-performance liquid chromatography 234
Saint Blanquat, G. De, see Alary, J. 485
Sakashita, T., see Nagata, O. 456
Saleh, S.
—, Johnston, A., Chanon, M. and Turner, P.
Determination of medifoxamine in plasma and urine by high-performance liquid chromatography 223
Sanders Krcmarik, C.
—, Miceli, J.J. and Pachla, L.A.
High-performance liquid chromatographic assay for a novel psychotropic drug (CI-943) in cynomolgus monkey plasma 493
Sauer, W.C., see Dugan, M.E.R. 430
Savory, J., see Herold, D.A. 180
Sawyer, R.C.
—, Stolfi, R.L. and Martin, D.S.
Quantitation of 5-fluorouracil incorporation into RNA by high-performance liquid chromatography without the use of radioactive precursors 450
Sekiya, T., see Matsui, H. 189
Shah, J.A.
— and Weber, D.J.
Simultaneous determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in urine by high-performance liquid chromatography with ultraviolet detection 245
Shimojo, M., see Shioya, H. 129
Shioya, H.
—, Shimojo, M. and Kawahara, Y.
Determination of a new angiotensin-converting enzyme inhibitor (CS-622) and its active metabolite in plasma and urine by gas chromatography–mass spectrometry using negative ion chemical ionization 129
Shukun, S.A.
— and Zav'yalov, V.P.
Peculiar features of application of pH gradients formed in borate buffer with a polyhydroxy compound for separation of protein in a free-flow electrophoretic apparatus 121
Siden, R., see Sintov, A. 335
Simpson, H., see Ashmore, S.P. 435
Singh, S., see Kinter, M. 201
Sintov, A.
—, Siden, R. and Levy, R.J.
Sensitive high-performance liquid chromatographic assay using 9-fluorenylmethylchloroformate for monitoring controlled-release lidocaine in plasma 335
Sion, J.-P.
—, Laureys, M., Gerlo, E. and Gorus, F.
Detection of macroenzymes in serum by high-performance gel permeation chromatography 91
Siu, T.-S.S., see Chan, Y.-P.M. 155
Smith, B.J., see Herold, D.A. 180
Smith, J.A.
— and O'Hare, M.
Effect of stationary and mobile phases on

hydrophobic interaction chromatography of proteins and peptides 71
Spahn, H., see Hartmann, C. 387
Stobaugh, J.F., see Mifune, M. 55
Stolfi, R.L., see Sawyer, R.C. 450
Svennerholm, L., see Davidsson, P. 279
Takahara, E., see Nagata, O. 456
Tan, Y., see Russel, F.G.M. 234
Tanaka, T.
—, Aramaki, S. and Momose, A.
Thin-layer chromatographic screening procedure for some drugs in horse plasma 407
Tanimura, T., see Nagao, T. 39
Thanh, H.H., see Rasmussen, K.E. 355
Thomson, A.H., see Ashmore, S.P. 435
Tønnesen, F., see Rasmussen, K.E. 355
Trechot, P., see Lapicque, F. 301
Tremaine, L.M.
— and Joerg, E.A.
Automated gas chromatographic-electron-capture assay for the selective serotonin uptake blocker sertraline 423
Turner, P., see Saleh, S. 223
Van Doorn, J.E.
—, Goormachtig, P.F.J. and De Haan, A.
Influence of perchloric acid on ion-pair high-performance liquid chromatography of nucleotides 441
Van Duyn, P., see Van Welie, R.T.H. 463
Van Ginneken, C.A.M., see Russel, F.G.M. 234
Van Meijel, J.J.M., see Russel, F.G.M. 234
Van Welie, R.T.H.
—, Van Duyn, P. and Vermeulen, N.P.E.
Determination of two mercapturic acid metabolites of 1,3-dichloropropene in human urine with gas chromatography and sulphur-selective detection 463
Vermeulen, N.P.E., see Baldew, G.S. 111
—, see Van Welie, R.T.H. 463
Virgili, P.
— and Henry, J.A.
Determination of lofepramine and desipramine using high-performance liquid chromatography and electrochemical detection 228
Wadgaonkar, N., see Ocampo, A.P. 167
Weber, D.J., see Shah, J.A. 245
Welie, R.T.H. van, see Van Welie, R.T.H. 463
Williams, E.D., see Brubaker, Jr., W.F. 377
Wills, M.R., see Herold, D.A. 180
Yamashita, H.
—, Kuroki, S. and Nakayama, F.
Assay of cholesterol 7α-hydroxylase utilizing a silica cartridge column and 5α-cholestane-3β,7β-diol as an internal standard 255
Yokoyama, Y., see Murata, K. 27
Yoshida, K., see Murata, K. 27
Zav'yalov, V.P., see Shukun, S.A. 121

Subject Index

N-Acetyl-S-(3-chloropropenyl-2)-L-cysteine
Determination of two mercapturic acid metabolites of 1,3-dichloropropene in human urine with GC and sulphur-selective detection 463

Ambenonium chloride
Sensitive determination of ambenonium chloride in serum from patients with myasthenia gravis using ion-exchange resin extraction and reversed-phase ion-pair chromatography 397

Amines
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39

Angiotensin-converting enzyme inhibitor
Determination of a new angiotensin-converting enzyme inhibitor (CS-622) and its active metabolite in plasma and urine by GC–MS using negative ion chemical ionization 129

Benzazepines
Quantitation of selective dopaminergic drugs in plasma by GC–MS following solid-phase extraction 201

Bilirubins
Review. Chromatographic analysis and structure determination of biliverdins and bilirubins 1

Biliverdins
Review. Chromatographic analysis and structure determination of biliverdins and bilirubins 1

Biogenic amines
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39

2,3-Butanediol isomers
GC determination of 2,3-butanediol isomers in urine 209

Carbamazepine
Simultaneous quantitation of salivary carbamazepine, carbamazepine-10,11-epoxide, phenytoin and phenobarbitone by HPLC 147

Carbamazepine-10,11-epoxide
Simultaneous quantitation of salivary carbamazepine, carbamazepine-10,11-epoxide, phenytoin and phenobarbitone by HPLC 147

2-Carboxythiazolidine-4-carboxylic acid
Reversed-phase ion-pair HPLC determination of 2-carboxythiazolidine-4-carboxylic acid in plasma 485

Celiprolol
Simultaneous determination of (*R*)- and (*S*)-celiprolol in human plasma and urine: HPLC assay on a chiral stationary phase with fluorimetric detection 387

9-Chloro-2-(2-furyl) [1,2,4] triazolo [1,5-*c*]-quinazolin-5-imine
Rapid, high-sensitivity reversed-phase HPLC assay for 9-chloro-2-(2-furyl)-[1,2,4]triazolo[1,5-*c*]quinazolin-5-imine and its oxo metabolite in plasma using fluorescence detection 377

Cholesterol 7α-hydroxylase
Assay of cholesterol 7α-hydroxylase utilizing a silica cartridge column and 5α-cholestane-3β,7β-diol as an internal standard 255

CI-943
HPLC assay for a novel psychotropic drug (CI-943) in cynomolgus monkey plasma 493

CS-622
Determination of a new angiotensin-converting enzyme inhibitor (CS-622) and its active metabolite in plasma and urine by GC–MS using negative ion chemical ionization 129

Desipramine
Determination of lofepramine and desipramine using HPLC and electrochemical detection 228

DL-Diaminopimelic acid
Cellulose clean-up and HPLC of DL-diaminopimelic acid in hydrolysates of physiological samples 430

3,4-Diaminopyridine
Analysis of 3,4-diaminopyridine in human serum by solid-phase extraction and HPLC with UV detection 214

1,3-Dichloropropene
Determination of two mercapturic acid metabolites of 1,3-dichloropropene in human urine with GC and sulphur-selective detection 463
Digoxin
Development of a HPLC–post-column fluorogenic assay for digoxin in serum 321
Dihydroxyvitamins D_3
TLC as an efficient alternative to HPLC in the assay of plasma vitamin D_3 metabolites 242
Dissaccharides
HPLC identification of disaccharides generated from heparan sulphate isomers using heparitinases 27
Dopamine
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39
Dopaminergic drugs
Quantitation of selective dopaminergic drugs in plasma by GC–MS following solid-phase extraction 201
Drugs
Quantitation of selective dopaminergic drugs in plasma by GC–MS following solid-phase extraction 201
Drugs
Determination of drug protein-binding by HPLC using a chemically bonded bovine albumin stationary phase 291
Drugs
TLC screening procedure for some drugs in horse plasma 407
Eicosanoids
15-Hydroxyeicosatetraenoic acid as a major eicosanoid in nasal secretions: assay by HPLC–radioimmunoassay and GC–MS procedures 416
Enkephalins
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Enzymes
Detection of macroenzymes in serum by high-performance gel permeation chromatography 91
Enzymes
Assay of ornithine decarboxylase activity by reversed-phase HPLC 101
Enzymes
Determination of a new angiotensin-converting enzyme inhibitor (CS-622) and its active metabolite in plasma and urine by GC–MS using negative ion chemical ionization 129
Enzymes
Assay of cholesterol 7α-hydroxylase utilizing a silica cartridge column and 5α-cholestane-3β,7β-diol as an internal standard 255
Epinepherine
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39
8-Ethyl-7,8-dihydro-1,3,5-trimethyl-1*H*-imidazo[1,2-*c*]pyrazolo[3,4-*e*]pyrimidine
HPLC assay for a novel psychotropic drug (CI-943) in cynomolgus monkey plasma 493
4′-Ethyl-2-methyl-3-(1-pyrrolidinyl)propiophenone hydrochloride
Determination of a new muscle relaxant (HY-770) in human serum by GC–MS 456
Exifone
Determination of exifone in human plasma and urine by HPLC with electrochemical detection 345
Flumequine
Solid-phase extraction and HPLC determination of flumequine and oxolinic acid in salmon plasma 355
9-Fluorenylmethylchloroformate
Sensitive HPLC assay using 9-fluorenylmethylchloroformate for monitoring controlled-release lidocaine in plasma 335
5-Fluorouracil
Quantitation of 5-fluorouracil incorporation into RNA by HPLC without the use of radioactive precursors 450
Furosemide
Solid-phase extraction of furosemide from plasma and urine and subsequent analysis by HPLC 234

Gangliosides
Gangliosides and sulphatide in human cerebrospinal fluid: quantitation with immunoaffinity techniques 279
Haemoglobins
Application of HPLC methodology to the analysis of hemoglobins synthesized in erythroid progenitor cells 83
Harmaline
Determination of pharmacological levels of harmane, harmine and harmaline in mammalian brain tissue, cerebrospinal fluid and plasma by HPLC with fluorimetric detection 269
Harmane
Determination of pharmacological levels of harmane, harmine and harmaline in mammalian brain tissue, cerebrospinal fluid and plasma by HPLC with fluorimetric detection 269
Harmine
Determination of pharmacological levels of harmane, harmine and harmaline in mammalian brain tissue, cerebrospinal fluid and plasma by HPLC with fluorimetric detection 269
Heparan sulphate isomers
HPLC identification of disaccharides generated from heparan sulphate isomers using heparitinases 27
Hippuric acid
HPLC separation of urinary hippuric and *o*-, *m*- and *p*-methylhippuric acids with a *β*-cyclodextrin-bonded column 189
Histamine
HPLC technique for the rapid determination of histamine in both the plasma and cellular components of blood 435
Homoharringtonine
Quantitation of homoharringtonine in plasma by HPLC with amperometric detection 155
HY-770
Determination of a new muscle relaxant (HY-770) in human serum by GC–MS 456
15-Hydroxyeicosatetraenoic acid
15-Hydroxyeicosatetraenoic acid as a major eicosanoid in nasal secretions: assay by HPLC–radioimmunoassay and GC–MS procedures 416
5-Hydroxytryptamine
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39
25-Hydroxyvitamin D_3
TLC as an efficient alternative to HPLC in the assay of plasma vitamin D_3 metabolites 242
Leucine-enkephalin
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Lidocaine
Sensitive HPLC assay using 9-fluorenylmethylchloroformate for monitoring controlled-release lidocaine in plasma 335
Lofepramine
Determination of lofepramine and desipramine using HPLC and electrochemical detection 228
Macroenzymes
Detection of macroenzymes in serum by high-performance gel permeation chromatography 91
Medifoxamine
Determination of medifoxamine in plasma and urine by HPLC 223
Methionine-enkephalin
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Methylhippuric acids
HPLC separation of urinary hippuric and *o*-, *m*- and *p*-methylhippuric acids with a *β*-cyclodextrin-bonded column 189
Methylprednisolone
Simultaneous determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human urine by HPLC with UV detection 245

Methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt
Simultaneous determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human urine by HPLC with UV detection 245
Metopimazine
HPLC method with fluorescence detection for the simultaneous determination of metopimazine and its acid metabolite in serum 472
β2-Microglobulin
Selective removal of β2-microglobulin from human plasma by high-performance immunoaffinity chromatography 194
Mobile phase composition
Effect of stationary and mobile phases on hydrophobic interaction chromatography of proteins and peptides 71
Muzolimine
Determination of muzolimine in plasma and urine by HPLC 478
Naphthalene-2,3-dicarboxaldehyde
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Non-steroidal anti-inflammatory drugs
Identification and simultaneous determination of non-steroidal anti-inflammatory drugs using HPLC 301
Norepinephrine
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39
Nucleotides
Influence of perchloric acid on ion-pair HPLC of nucleotides 441
Octopamine
Simultaneous determination of biogenic amines, their precursors and metabolites in a single brain of the cricket using HPLC with amperometric detection 39
Opioid peptides
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Ornithine decarboxylase
Assay of ornithine decarboxylase activity by reversed-phase HPLC 101
Oxolinic acid
Solid-phase extraction and HPLC determination of flumequine and oxolinic acid in salmon plasma 355
Peptides
Multi-dimensional HPLC of opioid peptides following pre-column derivatization with naphthalene-2,3-dicarboxaldehyde in the presence of cyanide ion. Preliminary results on the determination of leucine- and methionine-enkephalin-like fluorescence in the striatum region of the rat brain 55
Peptides
Effect of stationary and mobile phases on hydrophobic interaction chromatography of proteins and peptides 71
Phenobarbitone
Simultaneous quantitation of salivary carbamazepine, carbamazepine-10,11-epoxide, phenytoin and phenobarbitone by HPLC 147
Phenytoin
Simultaneous quantitation of salivary carbamazepine, carbamazepine-10,11-epoxide, phenytoin and phenobarbitone by HPLC 147
pH gradients
Peculiar features of application of pH gradients formed in borate buffer with a polyhydroxy compound for separation of proteins in a free-flow electrophoretic apparatus 121
Piperacillin
Determination of tazobactam and piperacillin in human plasma, serum, bile and urine by gradient elution reversed-phase HPLC 167

Piperazines
Determination of drug protein-binding by HPLC using a chemically bonded bovine albumin stationary phase 291
Polyhydroxy compound
Peculiar features of application of pH gradients formed in borate buffer with a polyhydroxy compound for separation of proteins in a free-flow electrophoretic apparatus 121
Protein-binding
Determination of drug protein-binding by HPLC using a chemically bonded bovine albumin stationary phase 291
Proteins
Effect of stationary and mobile phases on hydrophobic interaction chromatography of proteins and peptides 71
RS-5139
Determination of a new angiotensin-converting enzyme inhibitor (CS-622) and its active metabolite in plasma and urine by GC–MS using negative ion chemical ionization 129
Selenite
determination of ^{76}Se-labelled selenite and metabolites in plasma and urine by HPLC with on-line radioactivity detection 111
Serotonin uptake blocker
Automated GC–electron-capture assay for the selective serotonin uptake blocker sertraline 423
Sertraline
Automated GC–electron-capture assay for the selective serotonin uptake blocker sertraline 423
Stationary phases, hydrophobic interaction chromatography
Effect of stationary and mobile phases on hydrophobic interaction chromatography of proteins and peptides 71
Steroids
Simultaneous determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxo-octanoate] sodium salt in human urine by HPLC with UV detection 245
Sulphatide
Gangliosides and sulphatide in human cerebrospinal fluid: quantitation with immunoaffinity techniques 279
Sumatriptan
Development and validation of a LC–GC assay for the determination of sumatriptan in plasma 137
Tamoxifen
Determination of 2-methyl derivatives of tamoxifen in cell culture medium using HPLC and electrochemical detection 365
Tazobactam
Determination of tazobactam and piperacillin in human plasma, serum, bile and urine by gradient elution reversed-phase HPLC 167
Thromboxane B_2
Measurement of plasma thromboxane B_2 by GC–MS using $^{18}O_2$-labelled thromboxane B_2 as the internal standard 180
Vitamin D_3 metabolites
TLC as an efficient alternative to HPLC in the assay of plasma vitamin D_3 metabolites 242

Erratum

J. Chromatogr., 492 (1989) 137–165

Page 161, lines 15 and 16, "6-ketodinor-7,9,13-prostaglandin $F_{1\alpha}$" should read "6-ketoprostaglandin $F_{1\alpha}$".